



basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

SENIOR CERTIFICATE EXAMNATIONS ***SENIORSERTIFIKAAT-EKSAMEN***

MATHEMATICS P2/*WISKUNDE V2*

2017

MARKING GUIDELINES/*NASIENRIGLYNE*

MARKS: 150
PUNTE: 150

These marking guidelines consist of 22 pages.
Hierdie nasienriglyne bestaan uit 22 bladsye..

NOTE:

- If a candidate answers a question TWICE, only mark the FIRST attempt.
- If a candidate has crossed out an attempt of a question and not redone the question, mark the crossed out version.
- Consistent accuracy applies in ALL aspects of the marking memorandum. Stop marking at the second calculation error.
- Assuming answers/values in order to solve a problem is NOT acceptable.
- Geometry:
S = a mark for a correct statement (a statement mark is independent of a reason)
R = a mark for a correct reason (a reason mark may only be awarded if the statement is correct)
S/R = award a mark if statement and reason are both correct

NOTA:


- *As 'n kandidaat 'n vraag TWEEKEER beantwoord, merk slegs die EERSTE poging.*
- *As 'n kandidaat 'n antwoord van 'n vraag doodtrek en nie oordoen nie, merk die doodgetrekte poging.*
- *Volgehoue akkuraatheid word in ALLE aspekte van die memorandum toegepas. Hou op nasien by die tweede berekeningsfout.*
- *Aanvaar van antwoorde/waardes om 'n probleem op te los, word NIE toegelaat nie.*
- *Euklidiese Meetkunde:*
S = 'n punt vir 'n korrekte bewering ('n beweringspunt is onafhanklik van die rede)
R = 'n punt vir 'n korrekte rede ('n punt kan slegs vir 'n rede toegeken word, indien die bewering korrek is
S/R = 'n punt word toegeken indien beide die bewering en rede korrek is

QUESTION/VRAAG/VRAAG 1


TIME TAKEN (IN HOURS)	5	7	5	8	10	13	15	20	18	25	23
COST (IN THOUSANDS OF RANDS)	10	10	15	12	20	25	28	32	28	40	30

1.1	$a = 4,806... = 4,81$ $b = 1,323... = 1,32$ $y = 4,81 + 1,32x$	✓ $a = 4,81$ ✓ $b = 1,32$ ✓ equation (3)
1.2	Cost = 25,974... = 25,97 thousand rand (calculator) = R25 970 OR/OF $y = 4,81 + 1,32(16)$ $y = 25,93$ Cost = R25 930	✓ 25,97 ✓ answer (in Rands) (2) ✓ substitution ✓ answer (in Rands) (2)
1.3	$r = 0,949... = 0,95$	✓ answer (1)
1.4	$x = 0$ $y = 4,81$ OR (4,80647) \therefore R4 810 OR R4806,47	✓ $x = 0$ ✓ answer (2) [8]

QUESTION/VRAAG 2

2.1	modal class: $80 < x \leq 100$	✓ correct class (1)																		
2.2	<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <thead> <tr> <th style="padding: 5px;">Commission earned (in thousands of Rands)</th> <th style="padding: 5px;">Frequency</th> <th style="padding: 5px;">Cumulative Frequency</th> </tr> </thead> <tbody> <tr> <td style="padding: 5px;">$20 < x \leq 40$</td> <td style="padding: 5px;">7</td> <td style="padding: 5px;">7</td> </tr> <tr> <td style="padding: 5px;">$40 < x \leq 60$</td> <td style="padding: 5px;">6</td> <td style="padding: 5px;">13</td> </tr> <tr> <td style="padding: 5px;">$60 < x \leq 80$</td> <td style="padding: 5px;">8</td> <td style="padding: 5px;">21</td> </tr> <tr> <td style="padding: 5px;">$80 < x \leq 100$</td> <td style="padding: 5px;">10</td> <td style="padding: 5px;">31</td> </tr> <tr> <td style="padding: 5px;">$100 < x \leq 120$</td> <td style="padding: 5px;">4</td> <td style="padding: 5px;">35</td> </tr> </tbody> </table>	Commission earned (in thousands of Rands)	Frequency	Cumulative Frequency	$20 < x \leq 40$	7	7	$40 < x \leq 60$	6	13	$60 < x \leq 80$	8	21	$80 < x \leq 100$	10	31	$100 < x \leq 120$	4	35	<p>✓ 13 ; 21</p> <p>✓ 31 ; 35 (2)</p>
Commission earned (in thousands of Rands)	Frequency	Cumulative Frequency																		
$20 < x \leq 40$	7	7																		
$40 < x \leq 60$	6	13																		
$60 < x \leq 80$	8	21																		
$80 < x \leq 100$	10	31																		
$100 < x \leq 120$	4	35																		
2.3	<p style="text-align: center;">OGIVE</p> 	<p>✓ grounded/geanker</p> <p>✓ upper limits/ boonste limiet</p> <p>✓ cum frequency / Kum frekwensie</p> <p>✓ shape/vorm</p> <p style="text-align: right;">(4)</p>																		
2.4	<p>No. of salesmen awarded bonuses: $35 - 26$ = 9 salesmen</p>	<p>✓ accept (25 – 27)</p> <p>✓ accept (8 – 10)</p> <p style="text-align: right;">(2)</p>																		
2.5	<p>Estimated mean = $\frac{(30 \times 7) + (50 \times 6) + (70 \times 8) + (90 \times 10) + (110 \times 4)}{35}$</p> <p style="margin-left: 40px;">$= \frac{2410}{35}$</p> <p style="margin-left: 40px;">$= 68,86$ thousand rand or R68 857,14</p> <p style="margin-left: 40px;">$=$ R69 000 or 69 thousand rand</p>	<p>✓ top line using midpts & freq</p> <p>✓ 2410</p> <p>✓ answer (nearest)</p> <p style="text-align: right;">(3)</p> <p style="text-align: right;">[12]</p>																		


QUESTION/VRAAG 3



3.1	$m_{CD} = \frac{-3 - (-5)}{4 - 0}$ $= \frac{-3 + 5}{4 - 0}$ $= \frac{1}{2}$	✓ substitution of C & D ✓ answer (2)
3.2	$m_{AD} = \frac{-5 - 3}{0 - (-4)}$ $= -2$ $m_{CD} \times m_{AD} = \frac{1}{2} \times -2$ $= -1$ $\therefore AD \perp DC$	✓ substitution of A & D ✓ $m_{AD} = -2$ ✓ product = -1 (3)
3.3	$AB = \sqrt{(3 + 4)^2 + (4 - 3)^2} = \sqrt{50} = 5\sqrt{2}$ $BC = \sqrt{(4 - 3)^2 + (-3 - 4)^2} = 5\sqrt{2}$ $AB = BC$ $\therefore \Delta ABC \text{ is an isosceles triangle/'n gelykbenige driehoek}$	✓ correct substitution ✓ length of AB ✓ correct substitution ✓ length of BC (4)

<p>3.4</p>	$m_{CD} = m_{BF} = \frac{1}{2}$ $4 = \frac{1}{2}(3) + c$ $c = \frac{5}{2}$ $y = \frac{1}{2}x + \frac{5}{2}$ <p style="text-align: center;">OR/OF</p> $y - 4 = \frac{1}{2}(x - 3)$ $y - 4 = \frac{1}{2}x - 1\frac{1}{2}$ $y = \frac{1}{2}x + 2\frac{1}{2}$ <p style="text-align: center;">[BF DC]</p>	<p>✓ $m_{BF} = \frac{1}{2}$</p> <p>✓ substitution of B(3 ; 4)</p> <p>✓ equation (3)</p>
<p>3.5</p>	$\tan \alpha = -2$ $\therefore \alpha = 116,57^\circ$ $\alpha = 90^\circ + \theta$ <p style="text-align: center;">[ext $\angle \Delta$]</p> $\therefore \theta = 26,57^\circ$ <p>OR/OF</p> $\tan \alpha = -2 \text{ OR } m_{AD} = -2$ $\therefore \tan \theta = \frac{1}{2}$ $\therefore \theta = 26,57^\circ$ <p>OR/OF</p> <p>Inclination of DE is β:</p> $\tan \beta = \frac{1}{2}$ $\therefore \beta = 26,57^\circ$ $\therefore \hat{ODE} = 63,43^\circ$ $\therefore \theta = 90^\circ - 63,43^\circ$ $= 26,57^\circ$	<p>✓ $\tan \alpha = -2$</p> <p>✓ $\alpha = 116,57^\circ$</p> <p>✓ $\theta = 26,57^\circ$ (3)</p> <p>✓ $\tan \alpha = -2$</p> <p>✓ $\tan \theta = \frac{1}{2}$</p> <p>✓ $\theta = 26,57^\circ$ (3)</p> <p>✓ $\beta = 26,57^\circ$</p> <p>✓ $\hat{ODE} = 63,43^\circ$</p> <p>✓ $\theta = 26,57^\circ$ (3)</p>
<p>3.6</p>	$x^2 + y^2 = r^2$ $(4)^2 + (-3)^2 = 25$ $x^2 + y^2 = 25$	<p>✓ $r^2 = 25$</p> <p>✓ equation (2)</p> <p style="text-align: right;">[17]</p>

QUESTION/VRAAG 4




4.1	$N\left(\frac{1+(-3)}{2}; \frac{4+(-2)}{2}\right)$ <p>N(-1 ; 1) is the centre of the circle</p>	✓ substitution M & P ✓ x-value of N ✓ y-value of N (3)
4.2	$r = \sqrt{(1-(-1))^2 + (4-1)^2}$ $r = \sqrt{13} = \text{radius}$ $(x+1)^2 + (y-1)^2 = 13$ <p>OR/OR</p> $r = \sqrt{(-3-(-1))^2 + (-2-1)^2}$ $r = \sqrt{13} = \text{radius}$ $(x+1)^2 + (y-1)^2 = 13$	✓ substitution N & P ✓ $r = \sqrt{13}$ ✓ LHS of eq ✓ RHS of eq (4)
		✓ substitution N & M ✓ $r = \sqrt{13}$ ✓ LHS of eq ✓ RHS of eq (4)

<p>4.3</p>	<p>$m_{NM} \times m_{MR} = -1$ [radius \perp tangent/raakklyn] $m_{NM} = \frac{1 - (-2)}{-1 - (-3)}$ OR $m_{PM} = \frac{4 - (-2)}{1 - (-3)}$ $= \frac{3}{2}$ $= \frac{3}{2}$ $m_{MR} = -\frac{2}{3}$ $y - y_1 = -\frac{2}{3}(x - x_1)$ OR/OF $y = -\frac{2}{3}x + c$ $y + 2 = -\frac{2}{3}(x + 3)$ OR/OF $-2 = -\frac{2}{3}(-3) + c$ $y = -\frac{2}{3}x - 4$</p>	<p>✓ correct substitution ✓ m_{NM} ✓ m_{MR} ✓ substitution of m_{MR} & $(-3 ; -2)$ ✓ equation (5)</p>
<p>4.4</p>	<p>Symmetry of a kite: $S(-3 ; 4)$ OR/OF $\hat{P}SM = 90^\circ$ [\angle in semi circle] $PS \perp SM$ $\therefore S(-3 ; 4)$ OR/OF $(NS)^2 = (\text{radius})^2$ $(-3 + 1)^2 + (y - 1)^2 = 13$ $(y - 1)^2 = 9$ $y - 1 = \pm 3$ $y = 4$ OR $y \neq -2$ $\therefore S(-3 ; 4)$</p>	<p>✓ x-value of S ✓ y-value of S (2) ✓ x-value of S ✓ y-value of S (2) ✓ x-value of S ✓ y-value of S (2)</p>
<p>4.5</p>	<p>$(SR)^2 = (RM)^2$...Tangents from common pt/rklyne v dies punt $(x + 3)^2 + (y - 4)^2 = (x + 3)^2 + (y + 2)^2$ $y^2 - 8y + 16 = y^2 + 4y + 4$ $-12y = -12$ $y = 1$ $\frac{2}{3}x = -4 - 1$ or $1 = -\frac{2}{3}x - 4$ $x = -\frac{15}{2}$ $x = -7\frac{1}{2}$ $\therefore R\left(-7\frac{1}{2}; 1\right)$ OR/OF</p>	<p>✓ equating lengths ✓ simplification ✓ y-value of R ✓ x-value of R (4)</p>

	<p>$R(x;1)$ [RN is a horizontal line]</p> $\therefore 1 = -\frac{2}{3}x - 4$ $5 = -\frac{2}{3}x$ $x = -\frac{15}{2}$ $\therefore R\left(-\frac{15}{2};1\right)$ <p>OR/OF</p> $m_{NS} = \frac{1-4}{-1+3} = -\frac{3}{2}$ $\therefore m_{RS} = \frac{2}{3}$ $y-4 = \frac{2}{3}(x+3)$ $y = \frac{2}{3}x + 6$ $-\frac{2}{3}x - 4 = \frac{2}{3}x + 6$ $x = -7\frac{1}{2}$ $y = \frac{2}{3}\left(-\frac{15}{2}\right) + 6 = 1$ $\therefore R\left(-\frac{15}{2};1\right)$	<p>✓ $y_R = 1$ ✓ horizontal line OR R lies on $y = 1$ ✓ equating</p> <p>✓ x-value of R ($x < -4,6$)</p> <p>(4)</p> <p>✓ $y = \frac{2}{3}x + 6$</p> <p>✓ equating</p> <p>✓ x-value of R ($x < -4,6$)</p> <p>✓ y-value of R</p> <p>(4)</p>
<p>4.6</p>	<p>$RS = \sqrt{(-3+7,5)^2 + (4-1)^2}$ OR/OF $RM = \sqrt{(-3+7,5)^2 + (-2-1)^2}$</p> $RS = \frac{3\sqrt{13}}{2} = 5,41$ <p>area of RSNM = 2area of ΔRSN</p> $= 2\left(\frac{1}{2}\right)(\sqrt{13})\left(\frac{3\sqrt{13}}{2}\right)$ $= \frac{39}{2}$ <p>OR/OF 19,5 square units</p> <p>OR/OF</p>	<p>✓ RS OR RM</p> <p>✓ method</p> <p>✓ $\sqrt{13}$ and $\left(\frac{3\sqrt{13}}{2}\right)$</p> <p>✓ answer</p> <p>(4)</p> <p>✓ method</p> <p>✓ MS = 6</p> <p>✓ RN = 6,5</p> <p>✓ answer</p>

	<p>area RSNM = $\frac{1}{2}(MS \times RN)$ (area of a kite/<i>opp v vlieër</i>)</p> $= \frac{1}{2}(6)(6,5)$ $= \frac{39}{2} \text{ OR } 19,5 \text{ square units}$ <p>OR/OF</p> $RS = \sqrt{(-3 + 7,5)^2 + (4 - 1)^2} \text{ OR/OF } RM = \sqrt{(-3 + 7,5)^2 + (-2 - 1)^2}$ $RS = \frac{3\sqrt{13}}{2} \text{ or } 5,41$ $\text{area of } \Delta RSN = \left(\frac{1}{2}\right)(\sqrt{13})\left(\frac{3\sqrt{13}}{2}\right)$ $= \frac{39}{4} \text{ OR/OF } 9,75 \text{ square units}$ <p>area of RSNM = 2area of ΔRSN</p> $= \frac{39}{2} \text{ OR/OF } 19,5 \text{ square units}$ <p>OR/OF</p> <p>SM = 6</p> <p>area of RSNM = Area of ΔSMN + Area of ΔRSM</p> $= \frac{1}{2}(6)(1) + \frac{1}{2}(6)\left(5\frac{1}{2}\right)$ $= 3 + 16\frac{1}{2}$ $= 19\frac{1}{2}$	<p>(4)</p> <p>✓ RS OR RM</p> <p>✓ $\left(\frac{1}{2}\right)\sqrt{13}\left(\frac{3\sqrt{13}}{2}\right)$</p> <p>✓ method ✓ answer</p> <p>(4)</p> <p>✓ method</p> <p>✓ MS = 6</p> <p>✓ $h = 1$ & $5\frac{1}{2}$</p> <p>✓ answer</p> <p>(4)</p>
		<p>[22]</p>


QUESTION/VRAAG 5

<p>5.1.1</p>	$\tan A = \frac{\sin A}{\cos A}$ $= \frac{2p}{p}$ $= 2$ <p>OR/OF</p> $\tan A = \frac{2p}{p}$ $= 2$ <div style="text-align: center;">  <p>$(p ; 2p)$</p> </div>	<p>✓ identity</p> <p>✓ value of tan A (2)</p> <p>✓ $\frac{y}{x}$</p> <p>✓ value of tan A (2)</p>
<p>5.1.2</p>	$\sin^2 A + \cos^2 A = 1$ $(2p)^2 + p^2 = 1$ $4p^2 + p^2 = 1$ $5p^2 = 1$ $p^2 = \frac{1}{5}$ $\therefore p = -\frac{1}{\sqrt{5}}$	<p>✓ $(2p)^2 + p^2 = 1$</p> <p>✓ simplification of LHS</p> <p>✓ answer (3)</p>
<p>5.2</p>	$2 \sin^2 x - 5 \sin x + 2 = 0$ $(2 \sin x - 1)(\sin x - 2) = 0$ $\sin x = \frac{1}{2} \text{ or } \sin x = 2(\text{no solution})$ <p>ref $\angle = 30^\circ$</p> $\therefore x = 30^\circ + k.360^\circ \text{ or } x = 150^\circ + k.360^\circ ; k \in Z$	<p>✓ factors or formula</p> <p>✓ both equations</p> <p>✓ no solution/geen opl</p> <p>✓ $30^\circ + k.360^\circ$</p> <p>✓ $150^\circ + k.360^\circ ;$</p> <p>✓ $k \in Z$ (6)</p>
<p>5.3.1</p>	$\sin(x + 300^\circ) = \sin x \cos 300^\circ + \cos x \sin 300^\circ$	<p>✓ expansion/uitbreiding (1)</p>
<p>5.3.2</p>	$\sin(x + 300^\circ) - \cos(x - 150^\circ)$ $= \sin x \cos 300^\circ + \cos x \sin 300^\circ - (\cos x \cos 150^\circ + \sin x \sin 150^\circ)$ $= \sin x \cos 60^\circ - \cos x \sin 60^\circ - (-\cos x \cos 30^\circ + \sin x \sin 30^\circ)$ $= \sin x \cos 60^\circ - \cos x \sin 60^\circ + \cos x \cos 30^\circ - \sin x \sin 30^\circ$ $= \frac{1}{2} \sin x - \frac{\sqrt{3}}{2} \cos x + \frac{\sqrt{3}}{2} \cos x - \frac{1}{2} \sin x$ $= 0$ <p>OR/OF</p>	<p>✓ 2nd expansion/ 2de uitbreiding</p> <p>✓✓ reduction/reduksie</p> <p>✓ special angle values/ spesiale hoekwaardes</p> <p>✓ answer (5)</p>

	$\begin{aligned} & \sin(x + 300^\circ) - \cos(x - 150^\circ) \\ &= \sin x \cos 300^\circ + \cos x \sin 300^\circ - (\cos x \cos 150^\circ + \sin x \sin 150^\circ) \\ &= \sin x \cos 60^\circ - \cos x \sin 60^\circ - (-\cos x \cos 30^\circ + \sin x \sin 30^\circ) \\ &= \sin x \cos 60^\circ - \cos x \sin 60^\circ + \cos x \cos 30^\circ - \sin x \sin 30^\circ \\ &= \sin x \sin 30^\circ - \cos x \sin 60^\circ + \cos x \sin 60^\circ - \sin x \sin 30^\circ \\ &= 0 \end{aligned}$	<p>✓ 2nd expansion/ 2de uitbreiding ✓✓ reduction/reduksie ✓ co-ratios / ko-verh ✓ answer (5)</p>
<p>5.4</p>	<p>Consider: $\frac{\tan x + 1}{\sin x \tan x + \cos x} = \sin x + \cos x$</p> $\text{LHS} = \frac{\left(\frac{\sin x}{\cos x} + 1\right)}{\left(\sin x \cdot \frac{\sin x}{\cos x} + \cos x\right)} = \frac{\left(\frac{\sin x + \cos x}{\cos x}\right)}{\left(\frac{\sin^2 x + \cos^2 x}{\cos x}\right)}$ $= \frac{\sin x + \cos x}{\frac{\cos x}{1}}$ $= \frac{\sin x + \cos x}{\cos x} \times \frac{\cos x}{1}$ $= \sin x + \cos x$ <p>= RHS</p> <p>OR/OF</p> $\text{LHS} = \frac{\left(\frac{\sin x}{\cos x} + 1\right)}{\left(\sin x \cdot \frac{\sin x}{\cos x} + \cos x\right)} = \frac{\left(\frac{\sin x + \cos x}{\cos x}\right)}{\left(\frac{\sin^2 x + \cos^2 x}{\cos x}\right)}$ $= \frac{\left(\frac{\sin x}{\cos x} + 1\right)}{\frac{1}{\cos x}}$ $= \left(\frac{\sin x}{\cos x} + 1\right) \times \frac{\cos x}{1}$ $= \sin x + \cos x$ <p>= RHS</p>	<p>✓ identity of tan x ✓ $\frac{\sin x + \cos x}{\cos x}$ ✓ $\frac{\sin^2 x + \cos^2 x}{\cos x}$ ✓ $\sin^2 x + \cos^2 x = 1$ ✓ simplify (5)</p> <p>✓ identity of tan x ✓ $\frac{\sin^2 x + \cos^2 x}{\cos x}$ ✓ $\sin^2 x + \cos^2 x = 1$ ✓ simplify ✓ multiplication (5)</p>
<p>5.5.1</p>	$\begin{aligned} & (\sqrt{1+k})^2 = (\sin x + \cos x)^2 \\ & 1+k = \sin^2 x + 2 \sin x \cos x + \cos^2 x \\ & 1+k = 1 + \sin 2x \\ & k = \sin 2x \end{aligned}$	<p>✓ square both sides ✓ $\sin^2 x + \cos^2 x = 1$ ✓ sin 2x (3)</p>


5.5.2	<p>From 5.5.1</p> $\sin x + \cos x = \sqrt{1 + \sin 2x}$ $\therefore \text{max value: } \sin x + \cos x = \sqrt{1+1}$ $= \sqrt{2}$ <p>OR/OF</p> <p>Maximum value of $1 + \sin 2x = 1 + 1$</p> $= 2$ $\therefore \text{maximum value of } \sin x + \cos x = \sqrt{2}$ <p>OR/OF</p> $(\sin x + \cos x)^2 = \sin^2 x + 2 \sin x \cos x + \cos^2 x$ $= 1 + \sin 2x$ $\therefore \text{max value } (\sin x + \cos x)^2 = 1 + 1 = 2$ $\therefore \text{max value } \sin x + \cos x = \sqrt{2}$	<p>✓ max of $\sin 2x = 1$</p> <p>✓ answer (2)</p> <p>✓ max of $\sin 2x = 1$</p> <p>✓ answer (2)</p> <p>✓ max of $\sin 2x = 1$</p> <p>✓ answer (2)</p>
		[27]

QUESTION/VRAAG 6



6.1	Period = 180°	✓ answer (1)
6.2	-75°	✓ answer (1)
6.3	$\sin 2x \leq \frac{1}{\sqrt{2}} \cos x + \frac{1}{\sqrt{2}} \sin x$ $\sin 2x \leq \cos 45^\circ \cdot \cos x + \sin 45^\circ \cdot \sin x$ $\sin 2x \leq \cos(x - 45^\circ)$ $x \in [-75^\circ ; 165^\circ]$	✓ $\cos 45^\circ \cdot \cos x + \sin 45^\circ \cdot \sin x$ ✓ $\cos(x - 45^\circ)$ ✓ ✓ answer (4)
		[6]

QUESTION/VRAAG 7



Figure/Figuur (i)





Figure / Figuur (ii)

7.1	KC = 6 cm	✓ answer (1)
7.2	<p>Let P be the point of intersection of KL and CB</p> $\frac{KP}{KC} = \sin 60^\circ$ $KP = 6 \sin 60^\circ$ $KP = 3\sqrt{3} \text{ or } 5,20$ $\therefore KL = 8 + 3\sqrt{3} \text{ or } 13,20 \text{ cm}$	<p>✓ trig ratio</p> <p>✓ length of KP</p> <p>✓ answer (3)</p>
7.3	$DK^2 = 6^2 + 12^2$ $DK = \sqrt{180} \text{ or } 6\sqrt{5} \text{ or } 13,42 \text{ cm}$ $\frac{\sin \hat{KDL}}{KL} = \frac{\sin \hat{DLK}}{DK}$ $\frac{\sin \hat{KDL}}{\sin \hat{DLK}} = \frac{KL}{DK}$ $= \frac{8 + 3\sqrt{3}}{6\sqrt{5}} \text{ or } \frac{13,20}{13,42} \text{ or } 0,98$	<p>✓ DK = 6√5</p> <p>✓ use of sine rule</p> $\frac{\sin \hat{KDL}}{\sin \hat{DLK}} = \frac{KL}{DK}$ <p>✓ answer (4)</p> <p>[8]</p>


QUESTION/VRAAG 8



<p>8.1</p>	<p>$\hat{L} = 100^\circ$ [ext \angle cyclic quad = int opp \angle / <i>buite \angle kdvh = tos \angle</i>]</p> <p>OR/OF</p> <p>$\hat{N}_1 = 80^\circ$ [\angles on straight line]</p> <p>$\hat{L} = 100^\circ$ [opp \angles of cyclic quad]</p>	<p>✓S ✓R (2)</p> <p>✓S ✓R (2)</p>
<p>8.2</p>	<p>$\hat{N}_1 = 80^\circ$ [\angles on straight line/\anglee op reguitlyn]</p> <p>$\therefore \hat{O}_1 = 160^\circ$ [\angle at centre = $2 \times \angle$ at circumference/midpts $\angle = 2$ omtreks \angle]</p> <p>OR/OF</p> <p>reflex $\hat{K}\hat{O}\hat{M} = 200^\circ$ [\angle at centre = $2 \times \angle$ at circumference/midpts $\angle = 2 \times$ omtreks \angle]</p> <p>$\therefore \hat{O}_1 = 160^\circ$ [\angles around a pt/ \anglee om 'n pt]</p>	<p>✓S ✓S ✓R (3)</p> <p>✓S ✓R ✓S (3)</p>
<p>8.3</p>	<p>$\hat{M}_1 = 360^\circ - (100^\circ + 55^\circ + 160^\circ)$ [sum \angles of quad/som \anglee v vierhoek]</p> <p>$\therefore \hat{M}_1 = 45^\circ$</p>	<p>✓S ✓S (2)</p>


[7]

QUESTION/VRAAG 9



9.1.1	\angle in semi-circle/ \angle in halfsirkel	✓ answer (1)
9.1.2	Opp \angle s of quad = 180° / <i>Teenoorst</i> \angle e v vierhoek = 180°	✓ answer (1)
9.2.1	OF \perp AC [line from centre bisects chord/ <i>lyn v midpt halv kd</i>] \therefore AC \parallel GO [co-interior/ <i>ko-binne</i> \angle s = 180° OR/OF corresp/ooreenkomstige \angle s =]	✓ S ✓ R ✓ R (3)
9.2.2	$\hat{G}_1 = \hat{A}_2$ [corresp/ooreenk \angle s; AC \parallel GO] $\hat{A}_2 = \hat{B}_1$ [\angle s in same segment/ \angle e in dies segment] $\therefore \hat{G}_1 = \hat{B}_1$ OR/OF $\hat{G}_1 = \hat{B}_2$ [ext \angle cyclic quad/ <i>buite</i> \angle koordevh] but $\triangle ABF \cong \triangle CBF$ [s, \angle , s] $\therefore \hat{B}_2 = \hat{B}_1$ $\therefore \hat{G}_1 = \hat{B}_1$	✓ S ✓ R ✓ S ✓ R (4) ✓ S ✓ R ✓ R ✓ S (4)
9.3	OF : FB = 3 : 2 \therefore DO = 5k and DF = 8k $\therefore \frac{DG}{DA} = \frac{DO}{DF} = \frac{r}{\frac{8}{5}r}$ $\therefore \frac{DG}{DA} = \frac{5}{8}$	DB = 2r OR/OF DF = $2r - \frac{2}{5}r = \frac{8}{5}r$ [line \parallel side of Δ / <i>lyn</i> \parallel sy v Δ] ✓ S ✓ R ✓ S (3)
		[12]

QUESTION/VRAAG 10




10.1	Tangent-chord theorem	✓ R	(1)
10.2.1	$\hat{A}_2 + \hat{A}_3 = \hat{B}_1 + \hat{B}_2$ [∠ ^s opp = sides/∠ <i>eteenoor</i> = <i>sye</i>] $\hat{S}_3 = \hat{B}_1 + \hat{B}_2$ [ext ∠ cyclic quad/ <i>buite</i> ∠ <i>koordevh</i>] $\therefore \hat{S}_3 = \hat{A}_2 + \hat{A}_3$ $\therefore AB \parallel ST$ [corresp/ <i>ooreenk</i> ∠ ^s =]	✓ S ✓ R ✓ S ✓ R ✓ R	(5)
	OR/OF		
	$\hat{R}\hat{T}\hat{S} = \hat{B}\hat{A}\hat{S}$ [ext ∠ cyclic quad/ <i>buite</i> ∠ <i>koordevh</i>] $\hat{B}\hat{A}\hat{S} = \hat{A}\hat{B}\hat{T}$ [∠ ^s opp = sides/∠ <i>eteenoor</i> = <i>sye</i>] $\therefore \hat{R}\hat{T}\hat{S} = \hat{A}\hat{B}\hat{T}$ $\therefore AB \parallel ST$ [corresp/ <i>ooreenk</i> ∠ ^s =]	✓ S ✓ R ✓ S ✓ R ✓ R	(5)

10.2.2	$\hat{B}_2 = x$ [tan chord theorem/raakl – koordst] $x + \hat{T}_4 = \hat{B}_1 + \hat{B}_2$ [corresp/ooreenk \angle^s ; AB // ST] $\therefore \hat{T}_4 = \hat{B}_1$ $\hat{B}_1 = \hat{A}_1$ [tan chord theorem/raakl – koordst] $\therefore \hat{T}_4 = \hat{A}_1$	✓ S ✓ R ✓ S ✓ R ✓ R (5)
10.2.3	$\hat{T}_4 = \hat{A}_1$ [proven/bewys in 10.2.2] \therefore RTAP is a cyclic quadrilateral [line subtends = \angle^s] <i>Is 'n koordevierhoek [lyn onderspan = $\angle e$]</i>	✓ S ✓ R (2)
		[13]


QUESTION/VRAAG 11

11.1



<p>11.1</p>	<p>Constr: On sides AB and AC of $\triangle ABC$, mark points G and H respectively such that $AG = DE$ and $AH = DF$. Draw GH/Merk punt G en H op sy AB en AC van $\triangle ABC$ onderskeidelik af sodanig dat $AG = DE$ en $AH = DF$. Trek GH.</p> <p>Proof/Bewys:</p> <p>$\triangle AGH \equiv \triangle DEF$ [s, \angle, s]</p> <p>$\therefore \hat{A}GH = \hat{E}$</p> <p>$= \hat{B}$ [$\hat{B} = \hat{E}$, given/gegee]</p> <p>$\therefore GH \parallel BC$ [corresp/ooreenk $\angle^s =$]</p> <p>$\therefore \frac{AG}{AB} = \frac{AH}{AC}$ [line \parallel side of \triangle / lyn \parallel sye v \triangle]</p> <p>$\therefore \frac{DE}{AB} = \frac{DF}{AC}$ [constr/konstruksie]</p>	<p>✓ construction/ konstruksie</p> <p>✓ S/R</p> <p>✓ S</p> <p>✓ S /R</p> <p>✓ S ✓ R</p> <p>(6)</p>
-------------	---	--

11.2



<p>11.2.1(a)</p>	<p>$AP = PC$ [diag \parallel^m bisect each other/<i>hoekl \parallel^m halveer mekaar</i>] But $TP = PS$ [given/<i>gegee</i>] $AP - TP = PC - PS$ $\therefore AT = SC$</p>	<p>✓S ✓S OR S 2)</p>
<p>11.2.1(b)</p>	<p>In $\triangle PSR$ and $\triangle PBA$: $\hat{P}_1 = \hat{P}_3$ [vertically opp \angle^s / <i>regoorst $\angle e$</i>] $\hat{B}_1 = \hat{S}_1$ [\angle^s in same segment / <i>$\angle e$ in dies segment</i>] $\therefore \triangle PSR \parallel \triangle PBA$ [\angle, \angle, \angle]</p> <p>OR/OF In $\triangle PSR$ and $\triangle PBA$: $\hat{P}_1 = \hat{P}_3$ [vertically opp \angle^s / <i>regoorst $\angle e$</i>] $\hat{B}_1 = \hat{S}_1$ [\angle^s in same segment / <i>$\angle e$ in dies segment</i>] $\hat{A}_2 = \hat{R}_1$ [sum $\angle^s \Delta$ / <i>som $\angle e \Delta$</i>] $\therefore \triangle PSR \parallel \triangle PBA$ [\angle, \angle, \angle]</p>	<p>✓S ✓R ✓S ✓R ✓R (5)</p> <p>✓S ✓R ✓S ✓R ✓S (5)</p>

11.2.2(a)	$\frac{PR}{PA} = \frac{PS}{PB} \quad [\Delta s]$ $\therefore \frac{PR}{PA} = \frac{TR}{AD} = \frac{PS}{PB} \quad \left[\text{given } \frac{PR}{PA} = \frac{TR}{AD} \right]$ $\therefore \frac{PR}{PA} = \frac{TR}{AD} = \frac{TP}{PD} \quad [PS = TP; PB = PD]$ $\therefore \Delta RPT \parallel \Delta APD \quad [\text{sides of } \Delta \text{ in prop/sye v } \Delta \text{ in dies verhouding}]$	✓ S (all 3 ratios) ✓ S ✓ R (3)
11.2.2(b)	$\hat{T}_1 = \hat{D}_2 \quad [\Delta s]$ $\therefore \text{ATRD is a cyclic quad} \quad [\text{converse: ext } \angle \text{ of cyclic quad/ } \textit{Omgekeerde buite } \angle \text{ v koordevh}]$	✓ S ✓ R (2)
		[18]

TOTAL/TOTAAL: 150